

HAWKING RADIATION FROM BLACK HOLES OF CONSTANT NEGATIVE CURVATURE VIA GRAVITATIONAL ANOMALIES

PETROS SKAMAGOULIS

*Department of Physics, National Technical University of Athens,
Zografou Campus GR 157 73, Athens, Greece
E-mail: pskam@central.ntua.gr*

I derive the Hawking flux from black holes of constant negative curvature and from a black hole of constant negative curvature conformally coupled to a scalar field, using the covariant gravitational anomalies method.

1. Introduction

A few years ago Wilczek and collaborators proposed a new method to derive the Hawking flux from spherical black holes.^{1–3} They have proved, using a dimensional reduction procedure, that near the event horizon effective field theories become two-dimensional and exhibit gauge and gravitational anomalies. They showed, in the Unruh vacuum, that the Hawking flux emerges in order to cancel these anomalies. Recent applications and a modification of this method can be found in Refs. 4,5. A more technically simplified version of the method for asymptotically flat spacetimes, using only the covariant form of the anomalies, was suggested in Ref. 6. The purpose of this brief article, which is based in Ref. 7, is to apply this method of covariant anomalies in asymptotically non-flat spacetimes of non-spherical topology.

2. Hawking Radiation from Topological Black Holes

Consider the four-dimensional black hole solutions of Einstein's equations with negative cosmological constant $\Lambda = -3l^{-2}$. These asymptotically locally Anti-de Sitter (AdS) black holes are known as topological black holes (TBHs)⁸ and their metric is

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2(d\theta^2 + \sinh^2\theta d\varphi^2), \quad f(r) = -1 + \frac{r^2}{l^2} - \frac{2\mu}{r}, \quad (1)$$

where l is the AdS radius and $\mu > -l/3\sqrt{3}$ is a constant proportional to the black hole mass. Their topology is $\mathbb{R}^2 \times \Sigma$, where Σ is a two-dimensional manifold of constant negative curvature. The angular part of Eq. (1), with $0 \leq \varphi < 2\pi$ and $\theta \geq 0$, is the line element of the manifold Σ , which is locally isomorphic to the hyperbolic manifold H^2 and of the form $\Sigma = H^2/\Gamma$, where $\Gamma \subset O(2,1)$ is a discrete subgroup of isometries of H^2 . The manifold Σ is a compact two-dimensional surface of constant negative curvature and of genus $\bar{g} \geq 2$.⁷ Below, I will concentrate only on the genus $\bar{g} = 2$ case and deal with the event horizon r_H , which corresponds to the outer horizon for $-l/3\sqrt{3} < \mu < 0$ and to the unique horizon for $\mu \geq 0$.

Now consider a complex scalar field $\phi(x)$ in the background of the TBH of genus two. Its action is

$$S = -\frac{1}{2} \int d^4x \sqrt{-g} \phi^* \nabla^2 \phi + S_{\text{int}}, \quad (2)$$

where the first term is the free part of the action and S_{int} is the part which includes a mass term, potential terms and self-interaction terms. I perform a partial wave decomposition of ϕ in terms of the functions $\mathcal{Y}_\xi^m$, which are defined as

$$\mathcal{Y}_\xi^m(\theta, \varphi) = \left(\frac{2\pi}{\xi \tanh(\pi\xi)} \right)^{\frac{1}{2}} \frac{\Gamma(i\xi + \frac{1}{2})}{\Gamma(i\xi + m + \frac{1}{2})} P_{-\frac{1}{2}+i\xi}^m(\cosh \theta) e^{im\varphi}, \quad (3)$$

where $P_{-(1/2)+i\xi}^m$ are the associated Legendre functions, $m = 0, \pm 1, \pm 2, \pm 3, \dots$ and ξ takes discrete real values $\xi \geq 0$. These functions form a complete orthonormalized set of functions on the manifold Σ of genus two.^{7,9} Upon transforming to the “tortoise” coordinate r_* defined by $dr/dr_* = f(r)$ and taking the near-horizon limit $r_* \rightarrow -\infty$, one finds that the effective radial potentials for partial wave modes of the field, the mass term and the interaction terms contain the suppression factor $f(r(r_*))$ and vanish exponentially fast.⁷ Keeping only dominant terms, I find that physics near the event horizon can be effectively described by an infinite collection of two-dimensional free massless scalar fields, each propagating in a two-dimensional spacetime, which is given by the (t, r) part of the TBH metric of Eq. (1).⁷

In this reduced two-dimensional spacetime, following the ideas of Wilczek et al.,^{2,3} I neglect the ingoing modes in the region near the event horizon, which behave as left-moving modes, since they can not classically affect physics outside the horizon. Then, the effective two-dimensional theory exhibits a gravitational anomaly in this region, in the form of the non-conservation of the energy-momentum tensor.^{2,3} The time component of the covariant gravitational anomaly is^{6,7}

$$\partial_r \tilde{T}_t^r = \frac{1}{96\pi} f \partial_r f'' = \partial_r \tilde{N}_t^r, \quad \text{where} \quad \tilde{N}_t^r = \frac{1}{96\pi} \left(f f'' - \frac{f'^2}{2} \right), \quad (4)$$

$\tilde{T}_\nu^\mu$ is the covariant energy-momentum tensor and the prime denotes differentiation with respect to r . Solving this equation with the proposed in Refs. 2,3 boundary condition $\tilde{T}_t^r(r_H) = 0$, I find the non-conserved energy-momentum tensor

$$\tilde{T}_t^r(r) = \tilde{N}_t^r(r) - \tilde{N}_t^r(r_H). \quad (5)$$

The Hawking flux is measured at infinity, where there is no gravitational anomaly and in Refs. 1–3 it was given by the conserved energy-momentum tensor. One sees that the gravitational anomaly $\partial_r \tilde{N}_t^r$ vanishes at asymptotic infinity. However, $\tilde{N}_t^r$ is non-zero there, due to the fact that the spacetime asymptotically is AdS. These remarks and Eq. (4), which can be written as $\partial_r (\tilde{T}_t^r - \tilde{N}_t^r) = 0$, imply that the conserved energy-momentum tensor, or the energy flux F measured at infinity, is

$$F = \tilde{T}_t^r(r \rightarrow \infty) - \tilde{N}_t^r(r \rightarrow \infty) = -\tilde{N}_t^r(r_H) = \frac{\pi}{12} \left(\frac{f'(r_H)}{4\pi} \right)^2. \quad (6)$$

The energy flux F has a form equivalent to blackbody radiation of temperature $T = f'(r_H)/4\pi$,^{2,3} which is exactly the Hawking temperature of the TBH of Eq. (1).⁸ Hence, F can be identified with the Hawking flux from this TBH. However, I should note that the actual Hawking radiation observed at infinity is calculated through the grey-body factors, due to the curvature of the spacetime away from the horizon.¹

3. Hawking Radiation from a “Hairy” Topological Black Hole

Another interesting non-spherical black hole solution is a TBH conformally coupled to a scalar field Ψ . This black hole solution is known as the MTZ black hole¹⁰

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2 (d\theta^2 + \sinh^2 \theta d\varphi^2), \quad (7)$$

$$f(r) = \frac{r^2}{l^2} - \left(1 + \frac{G\mu}{r}\right)^2, \quad \Psi(r) = \sqrt{\frac{3}{4\pi G}} \frac{G\mu}{r + G\mu}, \quad (8)$$

where G is Newton’s constant and $\mu > -l/4G$ is a constant proportional to the black hole mass. The MTZ black hole has the same topology as the TBHs previously analyzed. Therefore, I will concentrate again only on the case of Σ being a genus two compact surface of constant negative curvature and deal with the event horizon r_+ , which is the unique horizon for $\mu \geq 0$ and the outer horizon for $-l/4 < G\mu < 0$.

Considering a complex scalar field ϕ in the background of the MTZ black hole of genus two, which does not interact with the scalar field Ψ , the dimensional reduction procedure proceeds as in Sec. 2. All the arguments of Sec. 2 hold also in this case⁷ and the conserved energy-momentum tensor is

$$F = \tilde{T}_t^r(r \rightarrow \infty) - \tilde{N}_t^r(r \rightarrow \infty) = -\tilde{N}_t^r(r_+) = \frac{\pi}{12} \left(\frac{f'(r_+)}{4\pi} \right)^2. \quad (9)$$

This energy flux has a form equivalent to blackbody radiation of temperature $T = f'(r_+)/4\pi$,^{2,3} which is the Hawking temperature of the MTZ black hole. The conformally coupled scalar field Ψ , which is time-independent, does not contribute to the Hawking flux. This was proved explicitly in Ref. 7 by showing that its action near the horizon vanishes, due to the presence of the suppression factor $f(r(r_*))$.

Conclusively, I derived the Hawking temperature and flux, without backscattering effects, from non-spherical black holes in asymptotically non-flat spacetimes, via the method of covariant anomalies, and showed that an infinite set of two-dimensional scalar fields near the horizon acts as the thermal source of this flux.

References

1. S. P. Robinson and F. Wilczek, *Phys. Rev. Lett.* **95**, 011303 (2005).
2. S. Iso, H. Umetsu and F. Wilczek, *Phys. Rev. Lett.* **96**, 151302 (2006).
3. S. Iso, H. Umetsu and F. Wilczek, *Phys. Rev. D* **74**, 044017 (2006).
4. L. Bonora, M. Cvitan, S. Pallua and I. Smolic, *Phys. Rev. D* **80**, 084034 (2009); J. J. Peng and S. Q. Wu, arXiv:0906.5121 [hep-th]; S. W. Wei, R. Li, Y. X. Liu and J. R. Ren, *Eur. Phys. J. C* **65**, 281 (2010); R. Becar, P. Gonzalez, G. Pulgar and J. Saavedra, arXiv:0808.1735 [gr-qc].
5. T. Morita, *Phys. Lett. B* **677**, 88 (2009).
6. R. Banerjee, *Int. J. Mod. Phys. D* **17**, 2539 (2009).
7. E. Papantonopoulos and P. Skamagoulis, *Phys. Rev. D* **79**, 084022 (2009).
8. L. Vanzo, *Phys. Rev. D* **56**, 6475 (1997).
9. E. N. Argyres, C. G. Papadopoulos, E. Papantonopoulos and K. Tamvakis, *J. Phys. A* **22**, 3577 (1989).
10. C. Martinez, R. Troncoso and J. Zanelli, *Phys. Rev. D* **70**, 084035 (2004).